

# Iowa Influenza Surveillance Network (IISN)

# Influenza-like Illness (ILI) and Other Respiratory Viruses

## **Weekly Activity Report**



All data presented in this report are provisional and may change as additional reports are received



Iowa Influenza Geographic Spread						
No Activity	Sporadic	Local	Regional	Widespread		

Note: See CDC activity estimates for definition www.cdc.gov/flu/weekly/overview.htm

Quick Stats	
Percent of influenza rapid test positive	9% (93/1045)
Percent of RSV rapid tests positive	7% (12/179)
Influenza-associated hospitalizations	21/6487 inpatients surveyed
Percent of outpatient visits for ILI	0.92% (baseline 1.9%)
Percent school absence due to illness	2.3%
Number of schools with ≥10% absence due to illness	1
Influenza-associated mortality -all ages (Cumulative)	1
Influenza-associated pediatric mortality (Cumulative)	0

Note: Deaths are considered influenza-associated when influenza is listed on the death certificate. This is an underestimate of influenza-related deaths. Cumulative mortality totals are from 10/1/2017 to the current week.

# Iowa statewide activity summary:


Influenza activity continues to increase in Iowa. The geographic spread of influenza in Iowa is sporadic. For this reporting week, the State Hygienic Laboratory confirmed 25 influenza A(H3) and 10 influenza B (Yamagata Lineage) viruses from submitted samples as well one influenza A and one influenza B positive specimens pending subtyping/lineage. Twenty-one influenza-related hospitalizations were reported from sentinel hospitals during this reporting week. The proportion of outpatient visits due to influenza-like illness (ILI) was 0.96 percent, which is below the regional baseline of 1.9. One long-term care facility reported an influenza outbreak this week. Surveillance sites reported detecting the following non-influenza respiratory illnesses with the most frequency: 87 rhinovirus/enterovirus, 36 parainfluenza virus type 1, 32 coronavirus, 22 RSV, and 11 adenovirus.

#### International activity summary - (WHO):

Influenza activity increased slightly in the temperate zone of the northern hemisphere while in the temperate zone of the southern hemisphere activity appeared to have decreased at inter-seasonal levels. In Central America and the Caribbean, influenza activity remained low. Worldwide, influenza A(H3N2) and B viruses accounted for the majority of influenza detections.

Visit <u>www.who.int/influenza/surveillance monitoring/updates/latest update GIP surveillance/en/</u> for more information. It was last updated 11/27/2017.

#### National activity summary - (CDC)-Last Updated in Week 48:



**Synopsis:** During week 48 (November 26-December 2, 2017), overall influenza activity increased slightly in the United States.

**Viral Surveillance**: The most frequently identified influenza virus type reported by public health laboratories during week 48 was influenza A. The percentage of respiratory specimens testing positive for influenza in clinical laboratories declined slightly.

**Pneumonia and Influenza Mortality:** The proportion of deaths attributed to pneumonia and influenza (P&I) was below the system-specific epidemic threshold in the National Center for Health Statistics (NCHS) Mortality Surveillance System.

**Influenza-associated Pediatric Deaths:** Two influenza-associated pediatric deaths were reported.

**Influenza-associated Hospitalizations:** A cumulative rate of 3.0 laboratory-confirmed influenza-associated hospitalizations per 100,000 population was reported.

**Outpatient Illness Surveillance:** The proportion of outpatient visits for influenza-like illness (ILI) was 2.3%, which is above the national baseline of 2.2%. Regions 1, 4, 6 and 7 reported ILI at or above region-specific baseline levels. Three states experienced high ILI activity; Puerto Rico and three states experienced moderate ILI activity; the District of Columbia and six states experienced low ILI activity; and New York City and 38 states experienced minimal ILI activity.

**Geographic Spread of Influenza:** The geographic spread of influenza in seven states was reported as widespread; Puerto Rico and 18 states reported regional activity; 18 states reported local activity; and the District of Columbia, the U.S. Virgin Islands and seven states reported sporadic activity; and Guam did not report.

Detailed information can be found online at www.cdc.gov/flu/weekly/.

#### **Laboratory surveillance program:**

The State Hygienic Laboratory (SHL) is the primary lab for influenza testing and reporting in Iowa. SHL reports the number of tests performed and the type and sub-type of positive tests to the influenza surveillance network daily.

Table 1: Influenza A viruses detected by SHL by age group


	CURRENT WEEK			CI	JMULATIVE (	10/1/17-0	CURRENT WE	EK)	
Age Group	A(H1N1) pdm09	A (H3)	Pending/ Not subtyped	Total	A(H1N1) pdm09	A (H1N1v)	A (H3)	Pending/ Not subtyped	Total
0-4	0	2	0	2 (8%)	6	0	12	1	19 (18%)
5-17	0	2	0	2 (8%)	0	0	14	0	14 (13%)
18-24	0	2	1	3 (12%)	0	0	7	1	8 (8%)
25-49	0	4	0	4 (15%)	1	1	11	0	13 (12%)
50-64	0	3	0	3 (12%)	0	0	13	0	13 (12%)
>64	0	12	0	12 (46%)	0	0	38	0	38 (36%)
Total	0 (0%)	25 (96%)	1 (4%)	26	7 (7%)	1 (1%)	95 (90%)	2 (2%)	105

Note: Cell counts of three or less are sometimes suppressed to protect confidentiality. Totals by age may not add up to totals by subtype/lineage due to missing age information. Only cases of lowa residents are included. Specimens listed as "not subtyped" were not able to be subtyped due to weak positive lab results. This can be due to poor collection, timing of collection or stage of infection.

Table 2: Influenza B viruses detected by SHL by age group

	CURRENT WEEK			CUMU	ILATIVE (10/1/1	L7- CURRENT	WEEK)	
Age Group	Victoria Lineage	Yamagata Lineage	Lineage Pending	Total	Victoria Lineage	Yamagata Lineage	Lineage Pending	Total
0-4	0	2	0	2 (18%)	0	2	0	2 (7%)
5-17	0	2	0	2 (18%)	0	5	0	5 (18%)
18-24	0	0	0	0 (0%)	0	3	0	3 (11%)
25-49	0	2	0	2 (18%)	0	6	0	6 (21%)
50-64	0	2	0	2 (18%)	0	6	0	6 (21%)
>64	0	2	1	3 (27%)	0	5	1	6 (21%)
Total	0 (0%)	10 (91%)	1 (9%)	11	0 (0%)	27 (96%)	1 (4%)	28

Note: Cell counts of three or less are sometimes suppressed to protect confidentiality. Totals by age may not add up to totals by subtype/lineage due to missing age information. Only cases of lowa residents are included.




#### Rapid influenza and RSV test surveillance:


The State Hygienic Laboratory (SHL) runs a weekly web-based survey program where laboratorians report the number of influenza and respiratory syncytial virus (RSV) rapid tests performed and the number of tests positive. This table includes only the number of patients tested for influenza or RSV at laboratory surveillance sites throughout the state. This table does not provide case counts.

Table 3: Percent of influenza rapid tests positive and number of tests performed by region for the present week

REGION*		RAPID IN	IFLUENZA TES	TS	S RAPID RSV TESTS		
REGION	Tested	Flu A	Flu B	% Positive	Tested	Positive	% Positive
Region 1 (Central)	267	12	7	7	22	3	14
Region 2 (NE)	55	1	0	2	11	3	27
Region 3 (NW)	141	9	2	8	77	3	4
Region 4 (SW)	59	10	0	17	5	1	20
Region 5 (SE)	26	0	2	8	11	0	0
Region 6 (Eastern)	497	41	9	10	53	2	4
Total	1045	73	20	9	179	12	7

Note: see map in the school section for a display of the counties in each region.







# Influenza-associated hospitalizations:

Sentinel hospitals that participate in IISN voluntarily track and report the number of influenza-associated hospitalizations and the total number of inpatients each week. Iowa hospitals interested in joining this surveillance program should contact Andy Weigel at 515-322-1937 or <a href="mailto:andy.weigel@idph.iowa.gov">andy.weigel@idph.iowa.gov</a> or more information.

Table 4: Number of influenza-associated hospitalization reported by age group

AGE	CURRENT WEEK	CUMULATIVE (10/1/17 – CURRENT WEEK)
Age 0-4	2	4
Age 5-24	2	4
Age 25-49	4	8
Age 50-64	4	11
Age >64	9	43
Total	21	70






#### Outpatient health care provider surveillance program (ILINet):

Outpatient health care providers who participate in the ILINet (a national influenza surveillance program) report the number of patients seen with influenza-like illness and the total number of patient visits each week. This system is a key part of Iowa's influenza surveillance. Iowa health care providers interested in joining this surveillance program should contact Andy Weigel at 515-322-1937 or andy.weigel@idph.iowa.gov for more information.

Table 5: Outpatient visits for influenza-like illness (ILI)


Week	Age 0-4	Age 5-24	Age 25-49	Age 50-64	Age > 64	Total ILI	ILI Percent
Week 46, ending November 18	7	8	1	0	1	17	0.45
Week 47, ending November 25	7	5	3	4	5	24	1.03
Week 48, ending December 2	8	7	10	2	9	36	0.92

Note: Influenza-like Illness is defined as a fever of ≥100° F as well as cough and/or sore throat.



# **Seasonal influenza vaccination:**

Seasonal influenza vaccination data in Iowa is based on doses reported to the Iowa Immunization Registry Information System (IRIS). IRIS is a confidential, computerized, population-based system that tracks immunization for children, adolescents and adults who are seen in a variety of public and private healthcare provider sites throughout the state of Iowa. For more information on the immunization data, contact Kim Tichy, IRIS coordinator at 515-281-4288 or <a href="mailto:kimberly.tichy@idph.iowa.gov">kimberly.tichy@idph.iowa.gov</a>.



Note: The data for the 2017-2018 season is provisional due to the lag between the vaccine administration date and the date reported to IRIS.

# **Long-term care outbreaks:**

Table 6: Number of long-term care outbreaks investigated

and or realised or resignation of the control of th						
REGION	CURRENT WEEK	CUMULATIVE (10/1/17 – CURRENT WEEK)				
Region 1 (Central)	0	0				
Region 2 (NE)	0	0				
Region 3 (NW)	0	0				
Region 4 (SW)	0	0				
Region 5 (SE)	1	1				
Region 6 (Eastern)	0	1				
Total	0	2				

Note: see map in the school section for a display of the counties in each region.

# School surveillance program

IDPH monitors illnesses in schools from two different types of reporting: 10 percent school absence reports and weekly sentinel illness reporting. Iowa schools (K-12) track and report (including non-influenza illnesses) when the number of students absent with illness reaches or exceeds 10 percent of total student enrollment. Iowa sentinel schools that participate in IISN voluntarily track and report absence due to all illness and the total enrollment each week. This data provides excellent trends for influenza activity as well as age-specific information used to target vaccination efforts and messages.





Table 7: Number of schools reporting >10% absenteeism due to any illness

REGION	CURRENT WEEK	CUMULATIVE (10/1/17 – CURRENT WEEK)
Region 1 (Central)	1	2
Region 2 (NE)	0	1
Region 3 (NW)	0	0
Region 4 (SW)	0	0
Region 5 (SE)	0	0
Region 6 (Eastern)	0	2
Total	0	4

Note: A regional map with the total of schools by county that report at least 10 percent of students absent due to illness for this current reporting week is displayed below (region numbers and abbreviations are displayed in large black font near the middle of each region).

# Number of schools with at least 10 percent of students absent, by county and region




### Non-influenza respiratory viruses:

The State Hygienic Laboratory (SHL) runs a weekly web-based survey program where laboratorians report the number of positive tests for non-influenza respiratory viruses. This table also includes the positive non-influenza virus tests reported from the Dunes Medical Laboratories at Mercy Medical Center in Sioux City. The table includes only the number of positive tests at laboratory surveillance sites throughout the state. The table does not provide case counts.

Table 8: Number of positive results for non-influenza respiratory virus collected by SHL, Mercy Dunes in Sioux City and Iowa Methodist Medical Center

Viruses	CURRENT WEEK	CUMULATIVE (10/1/17 – CURRENT WEEK)
Adenovirus	11	102
Parainfluenza Virus Type 1	36	327
Parainfluenza Virus Type 2	1	2
Parainfluenza Virus Type 3	1	16
Parainfluenza Virus Type 4	2	24
Rhinovirus/Enterovirus	87	777
Respiratory syncytial virus (RSV)	22	63
Human metapneumovirus (hMPV)	6	19
Coronavirus	32	111
Total	198	1441



# Other resources:

#### Vaccine:

Influenza vaccine recommendation: idph.iowa.gov/immtb/immunization/vaccine

CDC vaccine information: www.cdc.gov/flu/faq/flu-vaccine-types.htm

Vaccine finder: http://vaccinefinder.org/

#### Neighboring states' influenza information:

Illinois: www.dph.illinois.gov/topics-services/diseases-and-conditions/influenza/surveillance

Minnesota: www.health.state.mn.us/divs/idepc/diseases/flu/stats/index.html

Missouri: health.mo.gov/living/healthcondiseases/communicable/influenza/reports.php

South Dakota: doh.sd.gov/diseases/infectious/flu/

Wisconsin: flu.wisconsin.gov/